

L-QUADRI-ALGEBRAS

LIGONG LIU, XIANG NI, AND CHENGMING BAI*

ABSTRACT. Quadri-algebras introduced by Aguiar and Loday are a class of remarkable Loday algebras. In this paper, we introduce a notion of L-quadri-algebra with 4 operations satisfying certain generalized left-symmetry, as a Lie algebraic analogue of quadri-algebra such that the commutator of the sum of the 4 operations is a Lie algebra. Any quadri-algebra is an L-quadri-algebra. Moreover, L-quadri-algebras fit into the framework of the relationships between Loday algebras and their Lie algebraic analogues, extending the well known fact that the commutator of an associative algebra is a Lie algebra. We also give the close relationships between L-quadri-algebras and some interesting structures like Rota-Baxter operators, classical Yang-Baxter equation, some bilinear forms satisfying certain conditions.

1. INTRODUCTION

Quadri-algebras, which were introduced by Aguiar and Loday ([AL]), are a remarkable class of Loday algebras. The Loday algebras ([Va], or ABQR operad algebras in [EG]) refer to the series of algebras with a common property of “splitting associativity” ([Lo3]), which the first and maybe the most important class of Loday algebras, namely, the dendriform algebras, were introduced by Loday ([Lo1]) in 1995 with motivation from algebraic K -theory. The other Loday algebras include dendriform trialgebras ([E, LR3]), NS -algebras ([Le2]), octo-algebras ([Le3]), ennea-algebras ([Le1]) and dendriform-Nijenhuis algebras ([Le2]).

In fact, the Loday algebras were introduced due to their own interesting motivations and then developed as independent algebraic systems quickly. For example, dendriform algebras have been studied quite extensively with connections to operads ([Lo3]), homology ([Fr1]), Hopf algebras ([Cha, LR2]), Lie and Leibniz algebras ([Fr2]), combinatorics ([LR1]), arithmetic ([Lo2]) and quantum field theory ([Fo]) and so on (see [EMP] and the references therein). Quadri-algebras were introduced due to their close relationships with dendriform algebras in several ways like the relations between Rota-Baxter operators and infinitesimal bialgebras ([AL]). They are closely related to the study of ennea-algebras ([Le1]), operads ([EG]) and $\mathcal{O}$ -operators ([Bai3]).

In this paper, we introduce a notion of L-quadri-algebra due to several different motivations.

(1) Relationships with Loday algebras. An L-quadri-algebra is regarded as a Lie algebraic analogue of a quadri-algebra, which is one of the main motivations to introduce such a notion. It is known that the commutator of an associative algebra is a Lie algebra. It is

2000 *Mathematics Subject Classification.* 16W30, 17A30, 17B60.

Key words and phrases. Lie algebra, Loday algebra, quadri-algebra, classical Yang-Baxter equation, $\mathcal{O}$ -operator.

* Corresponding author.

quite interesting to find that certain commutator of a dendriform algebra is a pre-Lie algebra ([Cha, Ron]) which has already been introduced independently in geometry ([Ko, Vi]) and deformations of associative algebras ([G]) and then appear in many fields ([Bu] and the references therein). Moreover, a close relationship among Lie algebras, associative algebras, pre-Lie algebras and dendriform algebras is given as follows in the sense of commutative diagram of categories ([Cha]):

$$(1.1) \quad \begin{array}{ccc} \text{Lie algebra} & \leftarrow & \text{Pre-Lie algebra} \\ \uparrow & & \uparrow \\ \text{Associative algebra} & \leftarrow & \text{Dendriform algebra} \end{array}$$

Then it is natural to ask whether there are certain similar structures as of “Loday algebras” on Lie algebras. In fact, due to the different motivations, a notion of L-dendriform algebra was introduced in [BLN] as a Lie algebraic analogue of dendriform algebra, which extends the commutative diagram (1.1) to the following commutative diagram:

$$(1.2) \quad \begin{array}{ccccccc} \text{Lie algebra} & \leftarrow & \text{Pre-Lie algebra} & \leftarrow & \text{L-dendriform algebra} & & \\ & \swarrow & & \swarrow & & \swarrow & \\ & & \text{Associative algebra} & \leftarrow & \text{Dendriform algebra} & \leftarrow & \text{Quadri-algebra} \end{array}$$

As the above commutative diagram (1.2) at the level of associative algebras extends to the more Loday algebras (like octo-algebras), it is natural to find the corresponding algebraic structures at the level of Lie algebras. We will show that the L-quadri-algebras are chosen in a certain sense such that the following diagram including both diagrams (1.1) and (1.2) as sub-diagrams is commutative:

$$(1.3) \quad \begin{array}{ccccccccccc} \text{Lie algebra} & \leftarrow & \text{Pre-Lie algebra} & \leftarrow & \text{L-dendriform algebra} & \leftarrow & \text{L-quadri-algebra} & & & & \\ & \swarrow & & \swarrow & & \swarrow & & \swarrow & & \swarrow & \\ & & \text{Associative algebra} & \leftarrow & \text{Dendriform algebra} & \leftarrow & \text{Quadri-algebra} & \leftarrow & \text{Octo-algebra} \end{array}$$

(2) “Generalized left-symmetry”. A pre-Lie algebra $(A, \cdot)$ is also called a left-symmetric algebra since the associator

$$(1.4) \quad (x, y, z) = x \cdot (y \cdot z) - (x \cdot y) \cdot z, \quad \forall x, y, z \in A$$

is left-symmetric in x, y . Note that the associativity is as the same as the fact that the associator is zero. Furthermore, we show that for an L-dendriform algebra, there are the “generalized associators” with two operations satisfying certain “generalized left-symmetry” ([BLN]), whereas a dendriform algebra satisfies that the generalized associators are zero. In this sense, the so-called “common property of splitting associativity” ([Lo3]) for Loday algebras is understood as the generalized associators being zero, and it is reasonable to assume that their Lie algebraic analogues satisfy the generalized left-symmetry. It is true for L-quadri-algebras. That is, an L-quadri-algebra is an algebra with 4 operations satisfying that certain “generalized associators”

are “generalized left-symmetric”. As a direct consequence, the commutator of the sum of the 4 operations of an L-quadri-algebra is a Lie algebra and a quadri-algebra is a special L-dendriform algebra with the generalized associators being zero.

(3) Rota-Baxter operators. Rota-Baxter operators on associative algebras first arose in probability theory ([Bax]) and later became a subject in combinatorics ([Rot]) and a lot of fields in mathematics and physics ([E] and the references therein). A Lie algebraic version of Rota-Baxter operator is exactly the operator form of the famous classical Yang-Baxter equation (CYBE) ([Se]). As two commuting Rota-Baxter operators on an associative algebra or a Rota-Baxter operator on a dendriform algebra give a quadri-algebra, which is one of the main motivations to introduce the notion of quadri-algebra ([AL], there is a similar result for an L-quadri-algebra: three pairwise commuting Rota-Baxter operators on a Lie algebra or two commuting Rota-Baxter operators on a pre-Lie algebra or a Rota-Baxter operator on an L-dendriform algebra give an L-quadri-algebra. The above approach is another important motivation to introduce the notion of L-quadri-algebra, as it did for quadri-algebras ([AL]).

We briefly introduce the following two motivations involving the Lie algebras themselves.

(4) Some nondegenerate bilinear forms on L-dendriform algebras. L-quadri-algebras are regarded as the underlying algebraic structures of the L-dendriform algebras with a nondegenerate 2-cocycle, just as pre-Lie algebras are the underlying algebraic structures of the symplectic Lie algebras ([Chu]) and L-dendriform algebras are the underlying algebraic structures of the Hessian structures ([Sh]), that is, the pre-Lie algebras with a nondegenerate 2-cocycle.

(5) Classical Yang-Baxter equation and $\mathcal{O}$ -operators. In fact, pre-Lie algebras can be regarded as the algebraic structures behind the CYBE which plays an important role in integrable systems and quantum groups ([CP] and the references therein). It could be seen more clearly in terms of $\mathcal{O}$ -operators of a Lie algebra which was a generalization of (the operator form of) the CYBE in a Lie algebra ([Se, Ku]). Moreover, the CYBE and the Rota-Baxter operators on a Lie algebra are understood as the special $\mathcal{O}$ -operators which correspond to the co-adjoint representation and the adjoint representation respectively ([Bai1]). In terms of $\mathcal{O}$ -operators, there are analogues of the CYBE in pre-Lie algebras ([Bai2], S -equation) and L-dendriform algebras ([BLN], LD -equation). We will show that L-quadri-algebras are the algebraic underpinning of the LD -equation and $\mathcal{O}$ -operators on L-dendriform algebras. In fact, the axioms of an L-quadri-algebra are equivalent to the fact that the identity map is an $\mathcal{O}$ -operator (see Theorem 3.17 for details).

The paper is organized as follows. In Section 2, we recall some basic facts on pre-Lie algebras and L-dendriform algebras. In Section 3, we introduce the notion of L-quadri-algebra and then study some basic properties of L-quadri-algebras in terms of the $\mathcal{O}$ -operators of L-dendriform algebras. Their relationships with Loday algebras are also given. In the Appendix, in order to

be self-contained, we introduce a notion of L-octo-algebra and then give certain relationships between L-octo-algebras and L-quadri-algebras.

Throughout this paper, all algebras are finite-dimensional and over a field of characteristic zero. We also give some notations as follows. Let A be an algebra with an operation $*$.

(1) Let $L_*(x)$ and $R_*(x)$ denote the left and right multiplication operator respectively, that is, $L_*(x)y = R_*(y)x = x * y$ for any $x, y \in A$. We also simply denote them by $L(x)$ and $R(x)$ respectively without confusion. Moreover let $L_*, R_* : A \rightarrow gl(A)$ be two linear maps with $x \rightarrow L_*(x)$ and $x \rightarrow R_*(x)$ respectively. In particular, when A is a Lie algebra, we let $\text{ad}(x)$ denote the adjoint operator, that is, $\text{ad}(x)y = [x, y]$ for any $x, y \in A$.

(2) Let $r = \sum_i a_i \otimes b_i \in A \otimes A$. Set

$$(1.5) \quad r_{12} = \sum_i a_i \otimes b_i \otimes 1, r_{13} = \sum_i a_i \otimes 1 \otimes b_i, r_{23} = \sum_i 1 \otimes a_i \otimes b_i,$$

where 1 is a symbol which plays a similar role of the unit. The operation between two r s is defined in an obvious way. For example,

$$(1.6) \quad r_{12} * r_{13} = \sum_{i,j} a_i * a_j \otimes b_i \otimes b_j, r_{13} * r_{23} = \sum_{i,j} a_i \otimes a_j \otimes b_i * b_j, r_{23} * r_{12} = \sum_{i,j} a_j \otimes a_i * b_j \otimes b_i.$$

(3) Let V be a vector space. Let $\sigma : V \otimes V \rightarrow V \otimes V$ be the exchanging operator defined as

$$(1.7) \quad \sigma(x \otimes y) = y \otimes x, \forall x, y \in V.$$

On the other hand, any $r \in V \otimes V$ can be identified as a linear map from the dual space V^* to V in the following way:

$$(1.8) \quad \langle r(v^*), u^* \rangle = \langle u^* \otimes v^*, r \rangle, \forall u^*, v^* \in V^*,$$

where $\langle, \rangle$ is the ordinary pair between the vector space V and its dual space V^* . Moreover, any invertible linear map $T : V^* \rightarrow V$ induces a nondegenerate bilinear form $\mathcal{B}(,)$ on V by

$$(1.9) \quad \mathcal{B}(u, v) = \langle T^{-1}u, v \rangle, \quad \forall u, v \in V.$$

(4) Let V be a vector space. For any linear map $\rho : A \rightarrow gl(V)$, define a linear map $\rho^* : A \rightarrow gl(V^*)$ by

$$(1.10) \quad \langle \rho^*(x)v^*, u \rangle = -\langle v^*, \rho(x)u \rangle, \quad \forall x \in A, u \in V, v^* \in V^*.$$

2. PRE-LIE ALGEBRAS AND L-DENDRIFORM ALGEBRAS

2.1. Pre-Lie algebras and L-dendriform algebras.

Definition 2.1. Let A be a vector space with a bilinear product $(x, y) \rightarrow x \circ y$. A is called a *pre-Lie algebra* if for any $x, y, z \in A$, the associator

$$(2.1) \quad (x, y, z) = (x \circ y) \circ z - x \circ (y \circ z)$$

is symmetric in x, y , that is,

$$(2.2) \quad (x, y, z) = (y, x, z), \text{ or equivalently } (x \circ y) \circ z - x \circ (y \circ z) = (y \circ x) \circ z - y \circ (x \circ z), \forall x, y, z \in A.$$

Proposition 2.2. ([Bu]) *Let $(A, \circ)$ be a pre-Lie algebra.*

(1) *The commutator*

$$(2.3) \quad [x, y] = x \circ y - y \circ x, \quad \forall x, y \in A$$

defines a Lie algebra $\mathfrak{g}(A)$, which is called the sub-adjacent Lie algebra of A .

(2) *$L_\circ$ gives a representation of the Lie algebra $\mathfrak{g}(A)$, that is,*

$$(2.4) \quad L_\circ([x, y]) = L_\circ(x)L_\circ(y) - L_\circ(y)L_\circ(x), \quad \forall x, y \in A.$$

The following conclusion can be obtained directly from Definition 2.1 and Proposition 2.2.

Proposition 2.3. *Let $\mathfrak{g}$ be a vector space with a bilinear product $\circ$. Then $(\mathfrak{g}, \circ)$ is a pre-Lie algebra if and only if $(\mathfrak{g}, [,])$ defined by Eq. (2.3) is a Lie algebra and $(L_\circ, \mathfrak{g})$ is a representation of the Lie algebra $(\mathfrak{g}, [,])$.*

Definition 2.4. ([Lo1, BLN]) Let A be a vector space with two bilinear products denoted by $\triangleright$ and $\triangleleft : A \otimes A \rightarrow A$. $(A, \triangleright, \triangleleft)$ is called an *L-dendriform algebra* if for any $x, y, z \in A$,

$$(2.5) \quad x \triangleright (y \triangleright z) - (x \bullet y) \triangleright z = y \triangleright (x \triangleright z) - (y \bullet x) \triangleright z,$$

$$(2.6) \quad x \triangleright (y \triangleleft z) - (x \triangleright y) \triangleleft z = y \triangleleft (x \bullet z) - (y \triangleleft x) \triangleleft z.$$

where $x \bullet y = x \triangleright y + x \triangleleft y$. In particular, if both sides of Eqs. (2.5) and (2.6) are zero, then $(A, \triangleright, \triangleleft)$ is called a *dendriform algebra*.

Remark 2.5. Due to the left-symmetry of x, y in Eq. (2.5), in fact, there are 3 independent identities for a dendriform algebra, that is,

$$x \triangleright (y \triangleright z) = (x \bullet y) \triangleright z, \quad x \triangleright (y \triangleleft z) = (x \triangleright y) \triangleleft z, \quad y \triangleleft (x \bullet z) = (y \triangleleft x) \triangleleft z, \quad \forall x, y, z \in A.$$

Hence any dendriform algebra is an L-dendriform algebra.

Proposition 2.6. ([BLN]) *Let $(A, \triangleright, \triangleleft)$ be an L-dendriform algebra.*

(1) *If we define*

$$(2.7) \quad x \bullet y = x \triangleright y + x \triangleleft y, \forall x, y \in A,$$

then $(A, \bullet)$ is a pre-Lie algebra. $(A, \bullet)$ is called the associated horizontal pre-Lie algebra.

(2) *If we define*

$$(2.8) \quad x \circ y = x \triangleright y - y \triangleleft x, \forall x, y \in A,$$

then $(A, \circ)$ is a pre-Lie algebra. $(A, \circ)$ is called the associated vertical pre-Lie algebra.

(3) *The associated horizontal and vertical pre-Lie algebras $(A, \bullet)$ and $(A, \circ)$ of an L-dendriform algebra $(A, \triangleright, \triangleleft)$ have the same sub-adjacent Lie algebra $\mathfrak{g}(A)$ defined by*

$$(2.9) \quad [x, y] = x \triangleright y + x \triangleleft y - y \triangleright x - y \triangleleft x, \forall x, y \in A.$$

This Lie algebra is called the sub-adjacent Lie algebra of the L-dendriform algebra $(L, \triangleright, \triangleleft)$.

2.2. Bimodules and $\mathcal{O}$ -operators.

Definition 2.7. ([Bai2]) Let $(A, \circ)$ be a pre-Lie algebra and V be a vector space. Let $l, r : A \rightarrow gl(V)$ be two linear maps. (l, r, V) is called a *bimodule* of $(A, \circ)$ if

$$(2.10) \quad l(x)l(y) - l(x \circ y) = l(y)l(x) - l(y \circ x),$$

$$(2.11) \quad l(x)r(y) - r(y)l(x) = r(x \circ y) - r(y)r(x), \forall x, y \in A.$$

Proposition 2.8. ([BLN]) *Let A be a vector space with two bilinear products denoted by $\triangleright, \triangleleft : A \otimes A \rightarrow A$.*

(1) *$(A, \triangleright, \triangleleft)$ is an L-dendriform algebra if and only if $(A, \bullet)$ defined by Eq. (2.7) is a pre-Lie algebra and $(L_{\triangleright}, R_{\triangleleft}, A)$ is a bimodule.*

(2) *$(A, \triangleright, \triangleleft)$ is an L-dendriform algebra if and only if $(A, \circ)$ defined by Eq. (2.8) is a pre-Lie algebra and $(L_{\triangleright}, -L_{\triangleleft}, A)$ is a bimodule.*

Definition 2.9. ([BLN]) Let $(A, \triangleright, \triangleleft)$ be an L-dendriform algebra and V be a vector space. Let $l_{\triangleright}, r_{\triangleright}, l_{\triangleleft}, r_{\triangleleft} : A \rightarrow gl(V)$ be four linear maps. $(l_{\triangleright}, r_{\triangleright}, l_{\triangleleft}, r_{\triangleleft}, V)$ is called a *bimodule* of $(A, \triangleright, \triangleleft)$ if the following five equations hold (for any $x, y \in A$)

$$(2.12) \quad [l_{\triangleright}(x), l_{\triangleright}(y)] = l_{\triangleright}[x, y];$$

$$(2.13) \quad [l_{\triangleright}(x), l_{\triangleleft}(y)] = l_{\triangleleft}(x \circ y) + l_{\triangleleft}(y)l_{\triangleleft}(x);$$

$$(2.14) \quad r_{\triangleright}(x \triangleright y) = r_{\triangleright}(y)r_{\triangleright}(x) + r_{\triangleright}(y)r_{\triangleleft}(x) + [l_{\triangleright}(x), r_{\triangleright}(y)] - r_{\triangleright}(y)l_{\triangleleft}(x);$$

$$(2.15) \quad r_{\triangleright}(x \triangleleft y) = r_{\triangleleft}(y)r_{\triangleright}(x) + l_{\triangleleft}(x)r_{\triangleright}(y) + [l_{\triangleleft}(x), r_{\triangleleft}(y)];$$

$$(2.16) \quad [l_{\triangleright}(x), r_{\triangleleft}(y)] = r_{\triangleleft}(x \bullet y) - r_{\triangleleft}(y)r_{\triangleleft}(x),$$

where $x \circ y = x \triangleright y - y \triangleleft x$, and $x \bullet y = x \triangleright y + x \triangleleft y$.

In fact, according to [Sc], $(l_{\triangleright}, r_{\triangleright}, l_{\triangleleft}, r_{\triangleleft}, V)$ is a bimodule of an L-dendriform algebra $(A, \triangleright, \triangleleft)$ if and only if the direct sum $A \oplus V$ of the underlying vector spaces of A and V is turned into an L-dendriform algebra by defining multiplication in $A \oplus V$ by (for any $x, y \in A, u, v \in V$)

$$(2.17) \quad (x + u) \triangleright (y + v) = x \triangleright y + l_{\triangleright}(x)v + r_{\triangleright}(y)u,$$

$$(2.18) \quad (x + u) \triangleleft (y + v) = x \triangleleft y + l_{\triangleleft}(x)v + r_{\triangleleft}(y)u,$$

We denote it by $A \ltimes_{l_{\triangleright}, r_{\triangleright}, l_{\triangleleft}, r_{\triangleleft}} V$.

The following conclusion involves the notations given by Eq. (1.10).

Proposition 2.10. ([BLN]) *Let $(l_{\triangleright}, r_{\triangleright}, l_{\triangleleft}, r_{\triangleleft}, V)$ be a bimodule of an L-dendriform algebra $(A, \triangleright, \triangleleft)$. Then $(l_{\triangleright}^* + l_{\triangleleft}^* - r_{\triangleright}^* - r_{\triangleleft}^*, r_{\triangleright}^*, r_{\triangleleft}^* - l_{\triangleright}^*, -(r_{\triangleright}^* + r_{\triangleleft}^*), V^*)$ is a bimodule of $(A, \triangleright, \triangleleft)$.*

Definition 2.11. (1) ([Ku]) Let $\rho : \mathfrak{g} \rightarrow gl(V)$ be a representation of a Lie algebra $\mathfrak{g}$. A linear map $T : V \rightarrow \mathfrak{g}$ is called an $\mathcal{O}$ -operator of $\mathfrak{g}$ associated to (ρ, V) if T satisfies

$$(2.19) \quad [T(u), T(v)] = T(\rho(T(u))v - \rho(T(v))u), \forall u, v \in V.$$

In particular, an $\mathcal{O}$ -operator $R : \mathfrak{g} \rightarrow \mathfrak{g}$ of a Lie algebra $\mathfrak{g}$ associated to the adjoint representation $(\text{ad}, \mathfrak{g})$ is called a *Rota-Baxter operator of (weight zero) on $\mathfrak{g}$* , that is, R satisfies

$$(2.20) \quad [R(x), R(y)] = R([R(x), y] + [x, R(y)]), \quad \forall x, y \in \mathfrak{g}.$$

(2) ([BLN]) Let $(A, \circ)$ be a pre-Lie algebra and (l, r, V) be a bimodule. A linear map $T : V \rightarrow A$ is called an $\mathcal{O}$ -operator associated to (l, r, V) if T satisfies

$$(2.21) \quad T(u) \circ T(v) = T(l(T(u))v + r(T(v))u), \forall u, v \in V.$$

In particular, an $\mathcal{O}$ -operator of a pre-Lie algebra $(A, \circ)$ associated to the bimodule $(L_{\circ}, R_{\circ}, A)$ is called a *Rota-Baxter operator (of weight zero) on $(A, \circ)$* .

(3) ([BLN]) Let $(A, \triangleright, \triangleleft)$ be an L-dendriform algebra and $(l_{\triangleright}, r_{\triangleright}, l_{\triangleleft}, r_{\triangleleft}, V)$ be a bimodule. A linear map $T : V \rightarrow A$ is called an $\mathcal{O}$ -operator of $(A, \triangleright, \triangleleft)$ associated to $(l_{\triangleright}, r_{\triangleright}, l_{\triangleleft}, r_{\triangleleft}, V)$ if T satisfies (for any $u, v \in V$)

$$(2.22) \quad T(u) \triangleright T(v) = T[l_{\triangleright}(T(u))v + r_{\triangleright}(T(v))u],$$

$$(2.23) \quad T(u) \triangleleft T(v) = T[l_{\triangleleft}(T(u))v + r_{\triangleleft}(T(v))u].$$

In particular, an $\mathcal{O}$ -operator of an L-dendriform algebra $(A, \triangleright, \triangleleft)$ associated to the bimodule $(L_{\triangleright}, R_{\triangleright}, L_{\triangleleft}, R_{\triangleleft}, A)$ is called a *Rota-Baxter operator (of weight zero) on $(A, \triangleright, \triangleleft)$* .

Example 2.12. (1) ([Bai]) Let $T : V \rightarrow \mathfrak{g}$ be an $\mathcal{O}$ -operator of a Lie algebra $\mathfrak{g}$ associated to a representation (ρ, V) . Then there exists a pre-Lie algebra structure on V given by

$$(2.24) \quad u \circ v = \rho(T(u))v, \forall u, v \in V.$$

In particular, if $R : \mathfrak{g} \rightarrow \mathfrak{g}$ is a Rota-Baxter operator (of weight zero) on a Lie algebra $\mathfrak{g}$, then there exists a pre-Lie algebra structure on $\mathfrak{g}$ given by ([GS])

$$(2.25) \quad x \circ y = [R(x), y], \quad \forall x, y \in \mathfrak{g}.$$

(2) ([BLN]) Let $T : V \rightarrow A$ be an $\mathcal{O}$ -operator of a pre-Lie algebra $(A, \circ)$ associated to a bimodule (l, r, V) . Then there exists an L-dendriform algebra structure on V given by

$$(2.26) \quad u \triangleright v = l(T(u))v, u \triangleleft v = -r(T(u))v, \forall u, v \in V.$$

In particular, if $R : A \rightarrow A$ is a Rota-Baxter operator (of weight zero) on a pre-Lie algebra $(A, \circ)$, then there exists an L-dendriform algebra structure on A given by

$$(2.27) \quad x \triangleright y = R(x) \circ y, x \triangleleft y = -y \circ R(x), \forall x, y \in A.$$

(3) ([BLN]) Let $\mathfrak{g}$ be a Lie algebra and $\{R_1, R_2\}$ be a pair of commuting Rota-Baxter operators (of weight zero). Then there exists an L-dendriform algebra structure on $\mathfrak{g}$ given by

$$(2.28) \quad x \triangleright y = [R_1(R_2(x)), y], x \triangleleft y = [R_2(x), R_1(y)], \forall x, y \in \mathfrak{g}.$$

Remark 2.13. Let $\mathfrak{g}$ be a Lie algebra and $r \in \mathfrak{g} \otimes \mathfrak{g}$. Recall that r is a *solution of classical Yang-Baxter equation (CYBE) in $\mathfrak{g}$* if

$$(2.29) \quad [r_{12}, r_{13}] + [r_{12}, r_{23}] + [r_{13}, r_{23}] = 0.$$

When r is skew-symmetric, r is a solution of CYBE in $\mathfrak{g}$ if and only if r is an $\mathcal{O}$ -operator of $\mathfrak{g}$ associated to $(\text{ad}^*, \mathfrak{g}^*)$ ([Ku]). If, in addition, there exists a nondegenerate invariant symmetric bilinear form on $\mathfrak{g}$, then r is a solution of CYBE in $\mathfrak{g}$ if and only if r regarded as a linear map from $\mathfrak{g}$ to $\mathfrak{g}$ is a Rota-Baxter operator (of weight zero), that is r satisfies Eq. (2.20) ([Se]). In this sense, Eq. (2.20) is called the *operator form of the CYBE*.

Proposition 2.14. ([BLN]) Let $(A, \triangleright, \triangleleft)$ be an L-dendriform algebra and $r \in A \otimes A$ be skew-symmetric. Then r is an $\mathcal{O}$ -operator of $(A, \triangleright, \triangleleft)$ associated to $(L_\triangleright^* + L_\triangleleft^* - R_\triangleright^* - R_\triangleleft^*, R_\triangleright^*, R_\triangleleft^* - L_\triangleleft^*, -(R_\triangleright^* + R_\triangleleft^*), A^*)$ if and only if r satisfies the following equation:

$$(2.30) \quad r_{23} \triangleleft r_{13} = r_{13} \circ r_{12} + r_{23} \bullet r_{12}.$$

Moreover, if, in addition, r is invertible, then r satisfies Eq. (2.30) if and only if the (nondegenerate) bilinear form $\mathcal{B}$ induced by r through Eq. (1.9) satisfies

$$(2.31) \quad \mathcal{B}(x \triangleleft y, z) = -\mathcal{B}(y, z \circ x) + \mathcal{B}(x, z \bullet y), \quad \forall x, y, z \in A.$$

Definition 2.15. Let $(A, \triangleright, \triangleleft)$ be an L-dendriform algebra. Let $r \in A \otimes A$. Eq. (2.30) is called *LD-equation* in $(A, \triangleright, \triangleleft)$. On the other hand, a skew-symmetric bilinear form $\mathcal{B}$ on A satisfying Eq. (2.31) is called a *2-cocycle* of $(A, \triangleright, \triangleleft)$.

Remark 2.16. In fact, the *LD-equation* in an L-dendriform algebra is regarded as an analogue of the CYBE in a Lie algebra ([BLN], also see Remark 2.13).

3. L-QUADRI-ALGEBRAS.

3.1. Definition and basic properties.

Definition 3.1. Let A be a vector space with four bilinear products $\searrow, \nearrow, \nwarrow, \swarrow: A \otimes A \rightarrow A$. $(A, \searrow, \nearrow, \nwarrow, \swarrow)$ is called an *L-quadri-algebra* if for any $x, y, z \in A$,

$$(3.1) \quad x \searrow (y \searrow z) - (x * y) \searrow z = y \searrow (x \searrow z) - (y * x) \searrow z,$$

$$(3.2) \quad x \searrow (y \nearrow z) - (x \vee y) \nearrow z = y \nearrow (x \succ z) - (y \wedge x) \nearrow z,$$

$$(3.3) \quad x \searrow (y \nwarrow z) - (x \searrow y) \nwarrow z = y \nwarrow (x * z) - (y \nwarrow x) \nwarrow z,$$

$$(3.4) \quad x \nearrow (y \prec z) - (x \nearrow y) \nwarrow z = y \swarrow (x \wedge z) - (y \swarrow x) \nwarrow z,$$

$$(3.5) \quad x \searrow (y \swarrow z) - (x \succ y) \swarrow z = y \swarrow (x \vee z) - (y \prec x) \swarrow z,$$

where

$$(3.6) \quad x \succ y = x \searrow y + x \nearrow y, x \prec y = x \nwarrow y + x \swarrow y,$$

$$(3.7) \quad x \vee y = x \searrow y + x \swarrow y, x \wedge y = x \nearrow y + x \nwarrow y,$$

$$(3.8) \quad x * y = x \searrow y + x \nearrow y + x \nwarrow y + x \swarrow y = x \succ y + x \prec y = x \vee y + x \wedge y.$$

Remark 3.2. Let $(A, \searrow, \nearrow, \nwarrow, \swarrow)$ be an L-quadri-algebra. If $\nearrow = \nwarrow = \swarrow = 0$, then $(A, \searrow)$ is a pre-Lie algebra. If $\nwarrow = \searrow = \swarrow = 0$, or $\searrow = \nearrow = \swarrow = 0$, or $\nearrow = \nwarrow = \searrow = 0$ respectively, then $(A, \nearrow)$ or $(A, \nwarrow)$ or $(A, \swarrow)$ is an associative algebra respectively. For the cases that two operations among $\searrow, \nearrow, \nwarrow, \swarrow$ are zero, we have the following results:

- (i) If $\nearrow = \swarrow = 0$, then $(A, \searrow, \nwarrow)$ is an L-dendriform algebra.
- (ii) If $\nwarrow = \swarrow = 0$, then $(A, \searrow, \nearrow)$ is an L-dendriform algebra.
- (iii) If $\nearrow = \nwarrow = 0$, then $(A, \searrow, \swarrow)$ is an L-dendriform algebra.
- (iv) If $\searrow = \nearrow = 0$, then $(A, \swarrow, \nwarrow)$ is a dendriform algebra.
- (v) If $\searrow = \swarrow = 0$, then $(A, \nearrow, \nwarrow)$ is a dendriform algebra.
- (vi) If $\searrow = \nwarrow = 0$, then

$$x \nearrow (y \swarrow z) = y \swarrow (x \nearrow z),$$

$$y \nearrow (x \nearrow z) = (y \nearrow x - x \swarrow y) \nearrow z, \quad y \swarrow (x \swarrow z) = (y \swarrow x - x \nearrow y) \swarrow z.$$

Hence $(A, -, \nearrow, \swarrow^\sigma)$ is a dendriform algebra, where $x \swarrow^\sigma y = y \swarrow x$ for any $x, y \in A$.

Proposition 3.3. *Let $(A, \searrow, \nearrow, \nwarrow, \swarrow)$ be an L-quadri-algebra.*

(1) *The products given by Eq. (3.6) define an L-dendriform algebra $(A, \succ, \prec)$ which is called the associated vertical L-dendriform algebra of $(A, \searrow, \nearrow, \nwarrow, \swarrow)$*

(2) *The products given by Eq. (3.7) define an L-dendriform algebra $(A, \vee, \wedge)$ which is called the associated depth L-dendriform algebra of $(A, \searrow, \nearrow, \nwarrow, \swarrow)$.*

(3) *The products given*

$$(3.9) \quad x \triangleright y = x \searrow y - y \nwarrow x, \quad x \triangleleft y = x \nearrow y - y \swarrow x, \quad \forall x, y \in A,$$

define an L-dendriform algebra $(A, \triangleright, \triangleleft)$ which is called the associated horizontal L-dendriform algebra of $(A, \searrow, \nearrow, \nwarrow, \swarrow)$.

Proof. We only give an explicit proof of (3) since the proof of the other two cases is similar. In this case, we show that

$$\begin{aligned} x \triangleright (y \triangleright z) &= x \searrow (y \searrow z - z \nwarrow y) - (y \searrow z - z \nwarrow y) \nwarrow x, \\ (x \triangleright y) \triangleright z &= (x \searrow y - y \nwarrow x) \searrow z - z \nwarrow (x \searrow y - y \nwarrow x), \\ (x \triangleleft y) \triangleright z &= (x \nearrow y - y \swarrow x) \searrow z - z \nwarrow (x \nearrow y - y \swarrow x), \\ y \triangleright (x \triangleright z) &= y \searrow (x \searrow z - z \nwarrow x) - (x \searrow z - z \nwarrow x) \nwarrow y, \\ (y \triangleright x) \triangleright z &= (y \searrow x - x \nwarrow y) \searrow z - z \nwarrow (y \searrow x - x \nwarrow y), \\ (y \triangleleft x) \triangleright z &= (y \nearrow x - x \swarrow y) \searrow z - z \nwarrow (y \nearrow x - x \swarrow y). \end{aligned}$$

By the definition of L-quadri-algebra, we show that

$$(x \triangleright y) \triangleright z + (x \triangleleft y) \triangleright z + y \triangleright (x \triangleright z) - (y \triangleleft x) \triangleright z - (y \triangleright x) \triangleright z - x \triangleright (y \triangleright z) = 0.$$

Similarly we have

$$(x \triangleright y) \triangleleft z + y \triangleleft (x \triangleright z) + y \triangleleft (x \triangleleft z) - (y \triangleleft x) \triangleleft z - x \triangleright (y \triangleleft z) = 0.$$

Therefore $(A, \triangleright, \triangleleft)$ is an L-dendriform algebra. □

Corollary 3.4. *Let $(A, \searrow, \nearrow, \nwarrow, \swarrow)$ be an L-quadri-algebra and $(A, \triangleright, \triangleleft)$, $(A, \succ, \prec)$, $(A, \vee, \wedge)$ be the associated horizontal, vertical and depth L-dendriform algebras respectively. Let $x, y \in A$.*

(1) *If we define*

$$(3.10) \quad x \circ y = x \searrow y + x \swarrow y - y \nwarrow x - y \nearrow x = x \triangleright y - y \triangleleft x = x \vee y - y \wedge x,$$

then $(A, \circ)$ is the associated vertical pre-Lie algebra of both $(A, \triangleright, \triangleleft)$ and $(A, \vee, \wedge)$.

(2) If we define (also see Eq. (3.8))

$$(3.11) \quad x * y = x \searrow y + x \nearrow y + x \nwarrow y + x \swarrow y = x \succ y + x \prec y = x \vee y + x \wedge y,$$

then $(A, *)$ is the associated horizontal pre-Lie algebra of both $(A, \succ, \prec)$ and $(A, \vee, \wedge)$.

(3) If we define

$$(3.12) \quad x \bullet y = x \searrow y + x \nearrow y - y \nwarrow x - y \swarrow x = x \triangleright y + x \triangleleft y = x \succ y - y \prec x,$$

then $(A, \bullet)$ is the associated horizontal pre-Lie algebra of $(A, \triangleright, \triangleleft)$ and the associated vertical pre-Lie algebra of $(A, \succ, \prec)$.

(4) If we define

$$(3.13) \quad [x, y] = x \searrow y + x \swarrow y + x \nearrow y + x \nwarrow y - (y \searrow x + y \swarrow x + y \nearrow x + y \nwarrow x),$$

then $(A, [,])$ is a Lie algebra. $(A, [,])$ is called the sub-adjacent Lie algebra of $(A, \searrow, \nearrow, \nwarrow, \swarrow)$ and we denote it by $\mathfrak{g}(A)$.

Proof. It follows from Proposition 2.6 and Proposition 3.3. $\square$

Proposition 3.5. Let A be a vector space with four bilinear products denoted by $\searrow, \nearrow, \nwarrow, \swarrow: A \otimes A \rightarrow A$.

(1) $(A, \searrow, \nearrow, \nwarrow, \swarrow)$ is an L-quadri-algebra if and only if $(A, \triangleright, \triangleleft)$ defined by Eq. (3.9) is an L-dendriform algebra and $(L_{\searrow}, -L_{\nwarrow}, L_{\nearrow}, -L_{\swarrow}, A)$ is a bimodule.

(2) $(A, \searrow, \nearrow, \nwarrow, \swarrow)$ is an L-quadri-algebra if and only if $(A, \succ, \prec)$ defined by Eq. (3.6) is an L-dendriform algebra and $(L_{\searrow}, R_{\nearrow}, L_{\swarrow}, R_{\nwarrow}, A)$ is a bimodule.

(3) $(A, \searrow, \nearrow, \nwarrow, \swarrow)$ is an L-quadri-algebra if and only if $(A, \vee, \wedge)$ defined by Eq. (3.7) is an L-dendriform algebra and $(L_{\searrow}, R_{\swarrow}, L_{\nearrow}, R_{\nwarrow}, A)$ is a bimodule.

Proof. The conclusions can be obtained by a straightforward computation or a similar proof as of Proposition 3.11. $\square$

Remark 3.6. In the sense of the above conclusion (1), an L-quadri-algebra is understood as an algebra structure whose left multiplications give a bimodule structure on the underlying vector space of the L-dendriform algebra defined by certain commutators. It can be regarded as the “rule” of introducing the notion of L-quadri-algebra, which more generally, is the “rule” of introducing the notions of the Loday algebras and their Lie algebraic analogues ([Bai3], also see Proposition 2.3 and Proposition 2.8).

Corollary 3.7. Let $(A, \searrow, \nearrow, \nwarrow, \swarrow)$ be an L-quadri-algebra and $(A, \triangleright, \triangleleft)$, $(A, \succ, \prec)$, $(A, \vee, \wedge)$ be the associated horizontal, vertical and depth L-dendriform algebras of $(A, \searrow, \nearrow, \nwarrow, \swarrow)$ respectively.

(1) $(L_{\searrow}^*, -L_{\nwarrow}^*, -L_{\nearrow}^*, L_{\swarrow}^*, A^*)$ is a bimodule of $(A, \triangleright, \triangleleft)$.

- (2) $(L_{\circ}^*, R_{\nearrow}^*, L_{\triangleright}^*, -R_{\wedge}^*, A^*)$ is a bimodule of $(A, \succ, \prec)$.
 (3) $(L_{\bullet}^*, R_{\swarrow}^*, -L_{\triangleleft}^*, -R_{\searrow}^*, A^*)$ is a bimodule of $(A, \vee, \wedge)$.

Proof. It follows from Proposition 2.10 and Proposition 3.5. $\square$

Proposition 3.8. Let $(A, \searrow, \nearrow, \nwarrow, \swarrow)$ be an L-quadri-algebra.

(1) [**Transpose of an L-quadri-algebra**] Define four bilinear products $\searrow^t, \nearrow^t, \nwarrow^t, \swarrow^t$ on A by

$$(3.14) \quad x \searrow^t y = x \searrow y, \quad x \nwarrow^t y = x \nwarrow y, \quad x \nearrow^t y = x \swarrow y, \quad x \swarrow^t y = x \nearrow y, \quad \forall x, y \in A.$$

Then $(A, \searrow^t, \nearrow^t, \nwarrow^t, \swarrow^t)$ is an L-quadri-algebra which is called the transpose of $(A, \searrow, \nearrow, \nwarrow, \swarrow)$. Moreover (for any $x, y \in A$)

$$(3.15) \quad x \vee^t y = x \succ y, \quad x \wedge^t y = x \prec y, \quad x \triangleright^t y = x \triangleright y, \quad x \triangleleft^t y = -y \triangleleft x, \quad x \succ^t y = x \vee y, \quad x \prec^t y = x \wedge y.$$

(2) [**Symmetry Sym^a**] Define four bilinear products $\searrow^a, \nearrow^a, \nwarrow^a, \swarrow^a$ on A by

$$(3.16) \quad x \searrow^a y = x \searrow y, \quad x \nwarrow^a y = -y \nearrow x, \quad x \nearrow^a y = -y \nwarrow x, \quad x \swarrow^a y = x \swarrow y, \quad \forall x, y \in A.$$

Then $(A, \searrow^a, \nearrow^a, \nwarrow^a, \swarrow^a)$ is an L-quadri-algebra and (for any $x, y \in A$)

$$(3.17) \quad x \vee^a y = x \vee y, \quad x \wedge^a y = -y \wedge x, \quad x \triangleright^a y = x \succ y, \quad x \triangleleft^a y = -y \prec x, \quad x \succ^a y = x \triangleright y, \quad x \prec^a y = -y \triangleleft x.$$

(3) [**Symmetry Sym^b**] Define four bilinear products $\searrow^b, \nearrow^b, \nwarrow^b, \swarrow^b$ on A by

$$(3.18) \quad x \searrow^b y = x \searrow y, \quad x \nwarrow^b y = -y \nearrow x, \quad x \nearrow^b y = x \swarrow y, \quad x \swarrow^b y = -y \nwarrow x, \quad \forall x, y \in A.$$

Then $(A, \searrow^b, \nearrow^b, \nwarrow^b, \swarrow^b)$ is an L-quadri-algebra and (for any $x, y \in A$)

$$(3.19) \quad x \vee^b y = x \triangleright y, \quad x \wedge^b y = -y \triangleleft x, \quad x \triangleright^b y = x \succ y, \quad x \triangleleft^b y = x \prec y, \quad x \succ^b y = x \vee y, \quad x \prec^b y = -y \wedge x.$$

(3) [**Symmetry Sym^c**] Define four bilinear products $\searrow^c, \nearrow^c, \nwarrow^c, \swarrow^c$ on A by

$$(3.20) \quad x \searrow^c y = x \searrow y, \quad x \nwarrow^c y = -y \swarrow x, \quad x \nearrow^c y = -y \nwarrow x, \quad x \swarrow^c y = x \nearrow y, \quad \forall x, y \in A.$$

Then $(A, \searrow^c, \nearrow^c, \nwarrow^c, \swarrow^c)$ is an L-quadri-algebra and (for any $x, y \in A$)

$$(3.21) \quad x \vee^c y = x \succ y, \quad x \wedge^c y = -y \prec x, \quad x \triangleright^c y = x \vee y, \quad x \triangleleft^c y = -y \wedge x, \quad x \succ^c y = x \triangleright y, \quad x \prec^c y = x \triangleleft y.$$

(4) [**Symmetry Sym^d**] Define four bilinear products $\searrow^d, \nearrow^d, \nwarrow^d, \swarrow^d$ on A by

$$(3.22) \quad x \searrow^d y = x \searrow y, \quad x \nwarrow^d y = -y \swarrow x, \quad x \nearrow^d y = x \nearrow y, \quad x \swarrow^d y = -y \nwarrow x, \quad \forall x, y \in A.$$

Then $(A, \searrow^d, \nearrow^d, \nwarrow^d, \swarrow^d)$ is an L-quadri-algebra and (for any $x, y \in A$)

$$(3.23) \quad x \vee^d y = x \triangleright y, \quad x \wedge^d y = x \triangleleft y, \quad x \triangleright^d y = x \vee y, \quad x \triangleleft^d y = x \wedge y, \quad x \succ^d y = x \succ y, \quad x \prec^d y = -y \prec x.$$

Proof. In fact, there are 5 non-trivial symmetries in the axioms defining an L-quadri-algebra which are consistent with the symmetry group Σ_3 . By a straightforward checking, the above 5 cases correspond to these symmetries respectively. $\square$

Lemma 3.9. ([BLN]) *Let $(l_{\triangleright}, r_{\triangleright}, l_{\triangleleft}, r_{\triangleleft}, V)$ be a bimodule of an L -dendriform algebra $(A, \triangleright, \triangleleft)$.*

- (1) *$(l_{\triangleright}, -l_{\triangleleft}, V)$ is a bimodule of the associated vertical pre-Lie algebra $(A, \circ)$.*
- (2) *$(l_{\triangleright}, r_{\triangleleft}, V)$ is a bimodule of the associated horizontal pre-Lie algebra $(A, \bullet)$.*
- (3) *$(l_{\triangleright}, V)$ is a representation of the sub-adjacent Lie algebra $\mathfrak{g}(A)$.*

Corollary 3.10. *Let $(A, \searrow, \nearrow, \nwarrow, \swarrow)$ be an L -quadri-algebra.*

- (1) *$(L_{\searrow}, -L_{\nearrow}, A)$ is a bimodule of the associated pre-Lie algebra $(A, \circ)$.*
- (2) *$(L_{\searrow}, R_{\nwarrow}, A)$ is a bimodule of the associated pre-Lie algebra $(A, *)$.*
- (3) *$(L_{\searrow}, -L_{\swarrow}, A)$ is a bimodule of the associated pre-Lie algebra $(A, \bullet)$.*
- (4) *$(L_{\searrow}, A)$ is a representation of the sub-adjacent Lie algebra $\mathfrak{g}(A)$.*

Proof. It follows from Lemma 3.9, Corollary 3.4 and Proposition 3.5. $\square$

3.2. Constructions of L-quadri-algebras. For brevity, in this subsection, we only give the study involving the associated horizontal L -dendriform algebras. The corresponding study on the other associated L -dendriform algebras can be obtained by the transpose and symmetries of the L -dendriform algebras through Proposition 3.8.

Proposition 3.11. *Let $(l_{\triangleright}, r_{\triangleright}, l_{\triangleleft}, r_{\triangleleft}, V)$ be a bimodule of an L -dendriform algebra $(A, \triangleright, \triangleleft)$. If T is an $\mathcal{O}$ -operator associated to $(l_{\triangleright}, r_{\triangleright}, l_{\triangleleft}, r_{\triangleleft}, V)$, then there exists an L -quadri-algebra structure on V defined by (for any $u, v \in V$)*

$$(3.24) \quad u \searrow v = l_{\triangleright}(T(u))v, u \nwarrow v = -r_{\triangleright}(T(u))v, u \nearrow v = l_{\triangleleft}(T(u))v, u \swarrow v = -r_{\triangleleft}(T(u))v.$$

Therefore there is an L -dendriform algebra structure on V defined by Eq. (3.9) as the associated horizontal L -dendriform algebra of $(V, \searrow, \nearrow, \nwarrow, \swarrow)$ and T is a homomorphism of L -dendriform algebras. Furthermore, $T(V) = \{T(v) \mid v \in V\} \subset A$ is an L -dendriform subalgebra of $(A, \triangleright, \triangleleft)$ and there is an induced L -quadri-algebra structure on A given by

$$(3.25) \quad \begin{aligned} T(u) \searrow T(v) &= T(u \searrow v), T(u) \nearrow T(v) = T(u \nearrow v), \\ T(u) \nwarrow T(v) &= T(u \nwarrow v), T(u) \swarrow T(v) = T(u \swarrow v), \forall u, v \in V. \end{aligned}$$

Moreover, the corresponding associated horizontal L -dendriform algebra structure on $T(V)$ is just the L -dendriform algebra structure of $(A, \triangleright, \triangleleft)$ and T is a homomorphism of L -quadri-algebras.

Proof. By Eq. (3.24), for any $u, v, w \in V$, we show that

$$\begin{aligned} u \searrow (v \searrow w) - v \searrow (u \searrow w) &= [l_{\triangleright}(T(u)), l_{\triangleright}(T(v))]w = l_{\triangleright}[T(u), T(v)]w, \\ [u, v] \searrow w &= T(l_{\triangleright}(T(u))v + T(r_{\triangleright}(T(v))u) + T(l_{\triangleleft}(T(u))v + T(r_{\triangleleft}(T(v))u)) \\ &\quad - T(l_{\triangleright}(T(v))u + T(r_{\triangleright}(T(u))v) - T(l_{\triangleleft}(T(v))u + T(r_{\triangleleft}(T(u))v)) \\ &= l_{\triangleright}[T(u), T(v)]w. \end{aligned}$$

Therefore we show that $u \searrow (v \searrow w) - v \searrow (u \searrow w) = [u, v] \searrow w$. By a similar way, we show that Eqs. (3.2)-(3.5) holds for $(V, \searrow, \nearrow, \nwarrow, \swarrow)$. The other conclusions follow immediately. $\square$

Corollary 3.12. *Let R be a Rota-Baxter operator of (weight zero) on an L -dendriform algebra $(A, \triangleright, \triangleleft)$. Then there exists an L -quadri-algebra structure on A defined by*

$$(3.26) \quad x \searrow y = R(x) \triangleright y, x \nwarrow y = -y \triangleright R(x), x \nearrow y = R(x) \triangleleft y, x \swarrow y = -y \triangleleft R(x), \forall x, y \in A.$$

Its transpose is given by

$$(3.27) \quad x \searrow^t y = R(x) \triangleright y, x \nwarrow^t y = -y \triangleright R(x), x \nearrow^t y = -y \triangleleft R(x), x \swarrow^t y = R(x) \triangleleft y, \forall x, y \in A.$$

Its symmetries are given by (for any $x, y \in A$)

$$(3.28) \quad x \searrow^a y = R(x) \triangleright y, x \nwarrow^a y = -R(y) \triangleleft x, x \nearrow^a y = x \triangleright R(y), x \swarrow^a y = -y \triangleleft R(x);$$

$$(3.29) \quad x \searrow^b y = R(x) \triangleright y, x \nwarrow^b y = -R(y) \triangleleft x, x \nearrow^b y = -y \triangleleft R(x), x \swarrow^b y = x \triangleright R(y);$$

$$(3.30) \quad x \searrow^c y = R(x) \triangleright y, x \nwarrow^c y = x \triangleleft R(y), x \nearrow^c y = x \triangleright R(y), x \swarrow^c y = R(x) \triangleleft y;$$

$$(3.31) \quad x \searrow^d y = R(x) \triangleright y, x \nwarrow^d y = x \triangleleft R(y), x \nearrow^d y = R(x) \triangleleft y, x \swarrow^d y = x \triangleright R(y).$$

Proof. It follows from Proposition 3.11 by choosing the bimodule $(L_{\triangleright}, R_{\triangleright}, L_{\triangleleft}, R_{\triangleleft}, A)$. $\square$

Lemma 3.13. *Let $\{R_1, R_2\}$ be a pair of commuting Rota-Baxter operators (of weight zero) on a pre-Lie algebra $(A, \circ)$. Then R_2 is a Rota-Baxter operator (of weight zero) on the L -dendriform algebra defined by Eq. (2.27) with $R = R_1$.*

Proof. For any $x, y \in A$, we show that

$$\begin{aligned} R_2(x) \triangleright R_2(y) &= R_1(R_2(x)) \circ R_2(y) = R_2(R_1(x)) \circ R_2(y) \\ &= R_2(R_2 R_1(x) \circ y + R_1(x) \circ R_2(y)) = R_2(R_2(x) \triangleright y + x \triangleright R_2(y)); \\ R_2(x) \triangleleft R_2(y) &= -R_2(y) \circ R_1(R_2(x)) = -R_2(y) \circ R_2(R_1(x)) \\ &= -R_2(R_2(y) \circ R_1(x) + y \circ R_2 R_1(x)) = R_2(x \triangleleft R_2(y) + R_2(x) \triangleleft y). \end{aligned}$$

Note that the above identities involve the axioms of a Rota-Baxter operator (of weight zero) and $R_1 R_2 = R_2 R_1$. Hence R_2 is a Rota-Baxter operator (of weight zero) on $(A, \triangleright, \triangleleft)$. $\square$

Corollary 3.14. *Let $\{R_1, R_2\}$ be a pair of commuting Rota-Baxter operators (of weight zero) on a pre-Lie algebra $(A, \circ)$. Then there exists an L -quadri-algebra structure on A defined by (for any $x, y \in A$)*

$$(3.32) \quad x \searrow y = R_1 R_2(x) \circ y, x \nwarrow y = -R_1(y) \circ R_2(x), x \nearrow y = -y \circ R_1 R_2(x), x \swarrow y = R_2(x) \circ R_1(y).$$

Proof. It follows from Corollary 3.11 and Lemma 3.13. $\square$

By a similar proof as of Lemma 3.13, we have the following conclusion.

Lemma 3.15. *Let $\{R_1, R_2, R_3\}$ be three pairwise commuting Rota-Baxter operators (of weight zero) of a Lie algebra $\mathfrak{g}$. Then R_3 is a Rota-Baxter operator (of weight zero) on the L-dendriform algebra defined by Eq. (2.28).*

Corollary 3.16. *Let $\{R_1, R_2, R_3\}$ be three pairwise commuting Rota-Baxter operators of a Lie algebra $\mathfrak{g}$. Then there exists an L-quadri-algebra structure on $\mathfrak{g}$ defined by*

$$(3.33) \quad \begin{aligned} x \searrow y &= [R_1(R_2(R_3(x))), y], \quad x \swarrow y = [R_3(x), R_1(R_2(y))], \\ x \nearrow y &= [R_2(R_3(x)), R_1(y)], \quad x \nwarrow y = [R_1(R_3(x)), R_2(y)], \quad \forall x, y \in \mathfrak{g}. \end{aligned}$$

Proof. It follows from Corollary 3.12 and Lemma 3.15. $\square$

Theorem 3.17. *Let $(A, \triangleright, \triangleleft)$ be an L-dendriform algebra. Then there exists an L-quadri-algebra structure on A such that $(A, \triangleright, \triangleleft)$ is the associated horizontal L-dendriform algebra if and only if there exists an invertible $\mathcal{O}$ -operator of $(A, \triangleright, \triangleleft)$.*

Proof. Suppose that there exists an invertible $\mathcal{O}$ -operator T of $(A, \triangleright, \triangleleft)$ associated to a bimodule $(l_\triangleright, r_\triangleright, l_\triangleleft, r_\triangleleft, V)$. By Proposition 3.11, there exists an L-quadri-algebra structure on V given by Eq. (3.24). Therefore we define an L-quadri-algebra structure on A by Eq. (3.25) such that T is an isomorphism of L-quadri-algebras, that is,

$$\begin{aligned} x \searrow y &= T(l_\triangleright(x)T^{-1}(y)), \quad x \swarrow y = -T(r_\triangleright(x)T^{-1}(y)), \\ x \nearrow y &= T(l_\triangleleft(x)T^{-1}(y)), \quad x \nwarrow y = -T(r_\triangleleft(x)T^{-1}(y)), \quad \forall x, y \in A. \end{aligned}$$

Moreover, the associated horizontal L-dendriform algebra is $(A, \triangleright, \triangleleft)$ since (for any $x, y \in A$)

$$\begin{aligned} x \searrow y - y \swarrow x &= T(l_\triangleright(x)T^{-1}(y)) + T(r_\triangleright(y)T^{-1}(x)) = T(T^{-1}(x) \triangleright T^{-1}(y)) = x \triangleright y, \\ x \nearrow y - y \nwarrow x &= T(l_\triangleleft(x)T^{-1}(y)) + T(r_\triangleleft(y)T^{-1}(x)) = T(T^{-1}(x) \triangleleft T^{-1}(y)) = x \triangleleft y. \end{aligned}$$

Conversely, let $(A, \searrow, \nearrow, \swarrow, \nwarrow)$ be an L-quadri-algebra and $(A, \triangleright, \triangleleft)$ be the associated horizontal L-dendriform algebra. Then the identity map $\text{id} : A \rightarrow A$ is an $\mathcal{O}$ -operator of $(A, \triangleright, \triangleleft)$ associated to the bimodule $(L_\searrow, -L_\swarrow, L_\nearrow, -L_\nwarrow, A)$. $\square$

The following conclusion reveals the relationship between the L-dendriform algebras with a nondegenerate (skew-symmetric) 2-cocycle and L-quadri-algebras:

Proposition 3.18. *Let $(A, \triangleright, \triangleleft)$ be an L-dendriform algebra with a nondegenerate 2-cocycle $\mathcal{B}$. Then there exists an L-quadri-algebra structure $(A, \searrow, \nearrow, \swarrow, \nwarrow)$ on A defined by*

$$(3.34) \quad \begin{aligned} \mathcal{B}(x \searrow y, z) &= -\mathcal{B}(y, [x, z]), \quad \mathcal{B}(x \swarrow y, z) = \mathcal{B}(y, z \triangleright x), \\ \mathcal{B}(x \nearrow y, z) &= -\mathcal{B}(y, z \circ x), \quad \mathcal{B}(x \nwarrow y, z) = -\mathcal{B}(y, z \bullet x), \quad \forall x, y, z \in A, \end{aligned}$$

such that $(A, \triangleright, \triangleleft)$ is the associated horizontal L-dendriform algebra.

Proof. By Proposition 2.14, the invertible linear map $T : A^* \rightarrow A$ defined by Eq. (1.9) is an invertible $\mathcal{O}$ -operator of $(A, \triangleright, \triangleleft)$ associated to the bimodule $(L_{\triangleright}^* + L_{\triangleleft}^* - R_{\triangleright}^* - R_{\triangleleft}^*, R_{\triangleright}^*, R_{\triangleright}^* - L_{\triangleleft}^*, -(R_{\triangleright}^* + R_{\triangleleft}^*), A^*)$. By Theorem 3.17, there is an L-quadri-algebra structure on A defined by

$$\begin{aligned} \mathcal{B}(x \searrow y, z) &= \mathcal{B}(T(L_{\triangleright}^* + L_{\triangleleft}^* - R_{\triangleright}^* - R_{\triangleleft}^*)(x)T^{-1}(y)), z) = \langle (L_{\triangleright}^* + L_{\triangleleft}^* - R_{\triangleright}^* - R_{\triangleleft}^*)(x)T^{-1}(y), z \rangle \\ &= -\langle T^{-1}(y), [x, z] \rangle = -\mathcal{B}(y, [x, z]); \\ \mathcal{B}(x \swarrow y, z) &= \mathcal{B}(T(R_{\triangleright}^*)(x)T^{-1}(y)), z) = \langle R_{\triangleright}^*(x)T^{-1}(y), z \rangle = -\langle T^{-1}(y), z \triangleright x \rangle = -\mathcal{B}(y, z \triangleright x); \\ \mathcal{B}(x \nearrow y, z) &= \mathcal{B}(T(R_{\triangleright}^* - L_{\triangleleft}^*)(x)T^{-1}(y)), z) = -\langle T^{-1}(y), z \triangleright x - x \triangleleft z \rangle = -\mathcal{B}(y, z \circ x); \\ \mathcal{B}(x \swarrow y, z) &= \mathcal{B}(T(-R_{\triangleright}^* - R_{\triangleleft}^*)(x)T^{-1}(y)), z) = \langle T^{-1}(y), z \triangleright x + z \triangleleft x \rangle = \mathcal{B}(y, z \bullet x), \end{aligned}$$

for any $x, y, z \in A$, such that $(A, \triangleright, \triangleleft)$ is the associated horizontal L-dendriform algebra. $\square$

Remark 3.19. By Proposition 3.8, it is easy to get the transpose and symmetries of the L-quadri-algebras in Corollary 3.14, Corollary 3.16, Theorem 3.17, and Corollary 3.18, like Corollary 3.12.

Corollary 3.20. *Let $(A, \triangleright, \triangleleft)$ be an L-dendriform algebra with a nondegenerate 2-cocycle $\mathcal{B}$. Then there exist two other L-dendriform algebras $(A, \vee, \wedge)$ and $(A, \succ, \prec)$ given by*

$$(3.35) \quad \mathcal{B}(x \vee y, z) = -\mathcal{B}(y, x * z), \mathcal{B}(y \wedge x, z) = \mathcal{B}(x, y \triangleleft z), \forall x, y, z \in A,$$

$$(3.36) \quad \mathcal{B}(x \succ y, z) = -\mathcal{B}(y, x \circ z), \mathcal{B}(y \prec x, z) = -\mathcal{B}(x, z \triangleleft y), \forall x, y, z \in A,$$

respectively. Moreover, they are the associated depth and vertical L-dendriform algebras of the L-quadri-algebra $(A, \searrow, \nearrow, \swarrow, \nwarrow)$ given Eq. (3.34) respectively.

Proof. It follows from Proposition 3.3 and Proposition 3.18. $\square$

The following conclusion provides a construction of solutions of LD-equation in certain L-dendriform algebras from L-quadri-algebras:

Lemma 3.21. ([BLN]) *Let $(A, \triangleright, \triangleleft)$ be an L-dendriform algebra and $(l_{\triangleright}, r_{\triangleright}, l_{\triangleleft}, r_{\triangleleft}, V)$ be a bimodule. Let $(l_{\triangleright}^* + l_{\triangleleft}^* - r_{\triangleright}^* - r_{\triangleleft}^*, r_{\triangleright}^*, r_{\triangleright}^* - l_{\triangleleft}^*, -(r_{\triangleright}^* + r_{\triangleleft}^*), V^*)$ be the bimodule given in Proposition 2.10. Let $T : V \rightarrow A$ be a linear map which can be identified as an element in the vector space $(A \oplus V) \otimes (A \oplus V)$. Then $r = T - \sigma(T)$ is a skew-symmetric solution of LD-equation in the L-dendriform algebra $A \bowtie_{l_{\triangleright}^* + l_{\triangleleft}^* - r_{\triangleright}^* - r_{\triangleleft}^*, r_{\triangleright}^*, r_{\triangleright}^* - l_{\triangleleft}^*, -(r_{\triangleright}^* + r_{\triangleleft}^*)} V^*$ if and only if T is an $\mathcal{O}$ -operator of $(A, \triangleright, \triangleleft)$ associated to $(l_{\triangleright}, r_{\triangleright}, l_{\triangleleft}, r_{\triangleleft}, V)$.*

Corollary 3.22. *Let $(A, \searrow, \nearrow, \swarrow, \nwarrow)$ be an L-quadri-algebra and $(A, \triangleright, \triangleleft)$, $(A, \succ, \prec)$, $(A, \vee, \wedge)$ be the associated horizontal, vertical, and depth L-dendriform algebras respectively. Let $\{e_1, \dots,$*

$e_n\}$ be a basis of A and $\{e_1^*, \dots, e_n^*\}$ be the dual basis. Then

$$(3.37) \quad r = \sum_{i=1}^n (e_i \otimes e_i^* - e_i^* \otimes e_i)$$

is a skew-symmetric solution of LD-equation in the L-dendriform algebras

$$A \ltimes_{L_\bullet^*, -L_\blacktriangleleft^*, -L_\blacktriangleright^*, L_\blacktriangleright^*} A^*, \quad A \ltimes_{L_\circ^*, R_\blacktriangleright^*, L_\blacktriangleright^*, -R_\blacktriangleleft^*} A^*, \quad A \ltimes_{L_\bullet^*, R_\blacktriangleleft^*, -L_\blacktriangleright^*, -R_\blacktriangleleft^*} A^*,$$

respectively. Moreover there is a natural 2-cocycle $\mathcal{B}$ of these L-dendriform algebras induced by r through Eq. (1.9) which is given by

$$(3.38) \quad \mathcal{B}(x + a^*, y + b^*) = -\langle a^*, y \rangle + \langle x, b^* \rangle, \quad \forall x, y \in A, a^*, b^* \in A^*.$$

Proof. Since id is an $\mathcal{O}$ -operator of the L-dendriform algebra $(A, \triangleright, \triangleleft)$ associated to the bimodule $(L_{\searrow}, -L_{\nwarrow}, L_{\nearrow}, -L_{\swarrow}, A)$, the L-dendriform algebra $(A, \succ, \prec)$ associated to the bimodule $(L_{\searrow}, R_{\nearrow}, L_{\swarrow}, R_{\nwarrow}, A)$ and the L-dendriform algebra $(A, \vee, \wedge)$ associated to the bimodule $(L_{\searrow}, R_{\swarrow}, L_{\nearrow}, R_{\nwarrow}, A)$, the conclusion follows from Lemma 3.21. $\square$

3.3. Relationships with Loday algebras.

Definition 3.23. ([AL]) Let A be a vector space with four bilinear products $\searrow, \nearrow, \nwarrow, \swarrow: A \otimes A \rightarrow A$. $(A, \searrow, \nearrow, \nwarrow, \swarrow)$ is called a *quadri-algebra* if both of two sides of Eqs. (3.1)-(3.5) are zero.

Remark 3.24. Note that in the case of quadri-algebras, Eq. (3.1) only gives an independent identity due to the symmetry in x, y , whereas every other equation (Eqs. (3.2)-(3.5)) provides 2 independent identities. Therefore there are 9 independent identities for a quadri-algebra. As a direct consequence, any quadri-algebra is an L-quadri-algebra.

Remark 3.25. In fact, the both sides of Eqs. (2.5)-(2.6) and Eqs. (3.1)-(3.5) can be regarded as the “generalized associators”. In this sense, pre-Lie algebras, L-dendriform algebras and L-quadri-algebras satisfy that the associators are left-symmetric or the “generalized associators” are “generalized left-symmetric”, whereas associative algebras, dendriform algebras and quadri-algebras satisfy that the associators or the ‘generalized associators’ are zero respectively.

Proposition 3.26. Let $(A, \triangleright, \triangleleft)$ be a dendriform algebra.

- (1) ([Lo1]) The products given by Eq. (2.7) define an associative algebra $(A, \bullet)$.
- (2) ([Ron],[Cha]) The products given by Eq. (2.8) define a pre-Lie algebra $(A, \circ)$.

Proposition 3.27. Let $(A, \searrow, \nearrow, \nwarrow, \swarrow)$ be a quadri-algebra.

- (1) ([AL]) The products given by Eq. (3.6) define a dendriform algebra $(A, \succ, \prec)$.
- (2) ([AL]) The products given by Eq. (3.7) define a dendriform algebra $(A, \vee, \wedge)$.
- (3) ([BLN]) The products given by Eq. (3.9) define an L-dendriform algebra $(A, \triangleright, \triangleleft)$.

- (4) ([AL]) The products given by Eq. (3.8) define an associative algebra $(A, *)$.
- (5) ([BLN]) The products given by Eq. (3.10) define a pre-Lie algebra $(A, \circ)$.
- (6) ([BLN]) The products given by Eq. (3.12) define a pre-Lie algebra $(A, \bullet)$.
- (7) ([BLN]) The products given by Eq. (3.13) define a Lie algebra $(\mathfrak{g}(A), [,])$.

The notion of *octo-algebra* was introduced in [Le3] as a Loday algebra with 8 operations. In order to get the explicit relationships between octo-algebras and L-quadri-algebras, we need put both of them into a bigger framework. So we introduce a notion of *L-octo-algebra* as a natural generalization, which was put in the Appendix due to the complexity.

Definition 3.28. ([Le3]) With notations in the Appendix, let A be a vector space over with eight bilinear products denoted by $\searrow_1, \searrow_2, \nearrow_1, \nearrow_2, \nwarrow_1, \nwarrow_2, \swarrow_1, \swarrow_2: A \otimes A \longrightarrow A$. $(A, \searrow_1, \searrow_2, \nearrow_1, \nearrow_2, \nwarrow_1, \nwarrow_2, \swarrow_1, \swarrow_2)$ is called an *octo-algebra* if both sides of Eqs. (A1)-(A14) are zero.

Corollary 3.29. Any octo-algebra is an L-octo-algebra.

The following conclusion gives the explicit relationships between octo-algebras and L-quadri-algebras (see Proposition A2).

Proposition 3.30. With the notations in the Appendix. Let $(A, \searrow_1, \searrow_2, \nearrow_1, \nearrow_2, \nwarrow_1, \nwarrow_2, \swarrow_1, \swarrow_2)$ be an octo-algebra.

- (1) ([Le3]) $(A, \vee_2, \wedge_2, \wedge_1, \vee_1)$ is a quadri-algebra.
- (2) ([Le3]) $(A, \succ_2, \succ_1, \prec_1, \prec_2)$ is a quadri-algebra.
- (3) ([Le3]) $(A, \searrow_{12}, \nearrow_{12}, \nwarrow_{12}, \swarrow_{12})$ is a quadri-algebra.
- (4) $(A, \triangleright_1^2, \triangleleft_1^2, \triangleleft_2^1, \triangleright_2^1)$ is an L-quadri-algebra.

Summarizing the above study in this subsection, we have the following commutative diagram: (we omit “algebra” for abbreviation)

$$\begin{array}{ccccccc}
 \text{Lie} & \xleftarrow{\quad} & \text{Pre-Lie} & \xleftarrow{\quad}^+ & \text{L-dendriform} & \xleftarrow{\quad}^+ & \text{L-quadri} & \xleftarrow{\quad}^+ & \text{L-octo} \\
 & \nwarrow & \uparrow \in & \nwarrow & \uparrow \in & \nwarrow & \uparrow \in & \nwarrow & \uparrow \in \\
 & & \text{Associative} & \xleftarrow{\quad}^+ & \text{Dendriform} & \xleftarrow{\quad}^+ & \text{Quadri} & \xleftarrow{\quad}^+ & \text{Octo}
 \end{array}$$

where “ $\uparrow \in$ ” means the inclusion. “ $+$ ” means the operation $x \circ_1 y + x \circ_2 y$ and “ $-$ ” means the operation $x \circ_1 y - y \circ_2 x$.

3.4. Bilinear forms on L-quadri-algebras and LQ-equation.

Definition 3.31. Let $(A, \searrow, \nearrow, \nwarrow, \swarrow)$ be an L-quadri-algebra. A skew-symmetric bilinear form $\mathcal{B}$ on A is called *invariant* if $\mathcal{B}$ satisfies Eq. (3.34).

The following conclusion is obvious:

Proposition 3.32. *Let $(A, \nearrow, \nwarrow, \searrow, \swarrow)$ be an L-quadri-algebra and $\mathcal{B}$ be a skew-symmetric bilinear form. If $\mathcal{B}$ is invariant on A , then $\mathcal{B}$ is a 2-cocycle on the associated horizontal, vertical and depth L-dendriform algebras. Conversely, if $\mathcal{B}$ is a nondegenerate (skew-symmetric) 2-cocycle on an L-dendriform algebra $(A, \triangleright, \triangleleft)$, then $\mathcal{B}$ is invariant on the L-quadri-algebra given by Eq. (3.34).*

Next we consider the symmetric bilinear forms on an L-quadri-algebra $(A, \nearrow, \nwarrow, \searrow, \swarrow)$.

Theorem 3.33. *Let $(A, \searrow, \nearrow, \nwarrow, \swarrow)$ be an L-quadri-algebra and $r \in A \otimes A$ be symmetric. Let $(A, \triangleright, \triangleleft)$, $(A, \succ, \prec)$, $(A, \vee, \wedge)$ be the associated horizontal, vertical and depth L-dendriform algebras of $(A, \searrow, \nearrow, \nwarrow, \swarrow)$ respectively. Then the following conditions are equivalent:*

- (1) *r is an $\mathcal{O}$ -operator of $(A, \triangleright, \triangleleft)$ associated to the bimodule $(L_*, -L_{\nwarrow}^*, -L_{\wedge}^*, L_{\prec}^*, A^*)$.*
- (2) *r is an $\mathcal{O}$ -operator of $(A, \succ, \prec)$ associated to the bimodule $(L_{\circ}^*, R_{\nearrow}^*, L_{\triangleright}^*, -R_{\wedge}^*, A^*)$.*
- (3) *r is an $\mathcal{O}$ -operator of $(A, \vee, \wedge)$ associated to the bimodule $(L_{\bullet}^*, R_{\swarrow}^*, -L_{\triangleleft}^*, -R_{\prec}^*, A^*)$.*
- (4) *r satisfies*

$$(3.39) \quad r_{13} \triangleright r_{23} = -(r_{12} \vee r_{23} + r_{12} \wedge r_{23}) + r_{12} \nwarrow r_{13},$$

$$(3.40) \quad r_{13} \triangleleft r_{23} = (r_{12} \wedge r_{23}) - r_{12} \prec r_{13}.$$

Proof. We give the proof of the equivalence between (1) and (4). The proof of the other equivalences is similar (note that the symmetry of r is necessary). Let $\{e_1, \dots, e_n\}$ be a basis of A and $\{e_1^*, \dots, e_n^*\}$ be its dual basis. Suppose

$$e_i \searrow e_j = \sum_{k=1}^n a_{ij}^k e_k, \quad e_i \nwarrow e_j = \sum_{k=1}^n b_{ij}^k e_k, \quad e_i \nearrow e_j = \sum_{k=1}^n c_{ij}^k e_k, \quad e_i \swarrow e_j = \sum_{k=1}^n d_{ij}^k e_k,$$

$$r = \sum_{i,j=1}^n a_{ij} e_i \otimes e_j, \quad a_{ij} = a_{ji}.$$

So $r(e_i^*) = \sum_{j=1}^n a_{ij} e_j$. Note that

$$r(a^*) \triangleright r(b^*) = r[L_{\nwarrow}^*(r(a^*))b^* - L_{\nwarrow}^*(r(b^*))a^*],$$

is equivalent to (for any i, j, t)

$$\sum_{k,l=1}^n a_{ik} a_{jl} (a_{kl}^t - b_{lk}^t) = - \sum_{k,l=1}^n a_{ik} a_{lt} (a_{kl}^j + c_{kl}^j + b_{kl}^j + d_{kl}^j) + \sum_{k,l=1}^n a_{jk} a_{lt} b_{kl}^i,$$

which is precisely the coefficient of $e_i \otimes e_j \otimes e_t$ in

$$r_{13} \triangleright r_{23} = -(r_{12} \vee r_{23} + r_{12} \wedge r_{23}) + r_{12} \nwarrow r_{13}.$$

Similarly, we show that

$$r(a^*) \triangleleft r(b^*) = r[-L_{\wedge}^*(r(a^*))b^* + L_{\wedge}^*(r(b^*))a^*],$$

is equivalent to

$$r_{13} \triangleleft r_{23} = (r_{12} \wedge r_{23}) - r_{12} \prec r_{13}.$$

Hence the conclusion holds. $\square$

Theorem 3.34. *Let $(A, \searrow, \nearrow, \nwarrow, \swarrow)$ be an L-quadri-algebra and $r \in A \otimes A$. Suppose that r is symmetric and invertible. Then r satisfies Eqs. (3.39)-(3.40) if and only if the nondegenerate bilinear form $\mathcal{B}$ induced by r through Eq. (1.9) satisfies (for any $x, y, z \in A$)*

$$(3.41) \quad \mathcal{B}(x \swarrow y, z) = -\mathcal{B}(y, z \bullet x) - \mathcal{B}(x, z \vee y),$$

$$(3.42) \quad \mathcal{B}(x \nearrow y, z) = \mathcal{B}(y, x \wedge z - z \vee x) - \mathcal{B}(x, z \succ y).$$

Proof. Let $r = \sum_i a_i \otimes b_i$. Since r is symmetric, we have $\sum_i a_i \otimes b_i = \sum_i b_i \otimes a_i$. Therefore $r(v^*) = \sum_i \langle v^*, a_i \rangle b_i = \sum_i \langle v^*, b_i \rangle a_i$ for any $v^* \in A^*$. Since r is nondegenerate, for any $x, y, z \in A$, there exist $u^*, v^*, w^* \in A^*$ such that $x = r(u^*), y = r(v^*), z = r(w^*)$. So

$$\begin{aligned} \mathcal{B}(x \swarrow y, z) &= \langle r(u^*) \swarrow r(v^*), w^* \rangle = \sum_{i,j} \langle u^*, b_i \rangle \langle v^*, b_j \rangle \langle w^*, a_i \swarrow a_j \rangle = \langle w^* \otimes u^* \otimes v^*, r_{12} \swarrow r_{13} \rangle; \\ \mathcal{B}(z \bullet x, y) &= \langle r(w^*) \bullet r(u^*), v^* \rangle = \sum_{i,j} \langle w^*, b_i \rangle \langle u^*, b_j \rangle \langle v^*, a_i \bullet a_j \rangle = \langle w^* \otimes u^* \otimes v^*, r_{13} \bullet r_{23} \rangle; \\ \mathcal{B}(z \vee y, x) &= \langle r(w^*) \vee r(v^*), u^* \rangle = \sum_{i,j} \langle w^*, b_i \rangle \langle v^*, b_j \rangle \langle u^*, a_i \vee a_j \rangle = \langle w^* \otimes u^* \otimes v^*, r_{12} \vee r_{23} \rangle. \end{aligned}$$

Hence Eq. (3.41) holds if and only if

$$(3.43) \quad r_{13} \bullet r_{23} = -r_{12} \swarrow r_{13} - r_{12} \vee r_{23},$$

which is exactly the sum of Eqs. (3.39) and (3.40). Similarly, Eq. (3.40) holds if and only if

$$(3.44) \quad \mathcal{B}(x \triangleleft y, z) = \mathcal{B}(y, x \vee z) - \mathcal{B}(x, y \prec z), \quad \forall x, y, z \in A.$$

Note that the above equation is exactly the difference of Eqs. (3.41) and (3.42):

$$(3.45) \quad \mathcal{B}(x \nearrow y, z) - \mathcal{B}(y \swarrow x, z) = \mathcal{B}(y, x \wedge z - z \vee x) - \mathcal{B}(x, z \succ y) + \mathcal{B}(x, z \bullet y) + \mathcal{B}(y, z \vee x).$$

So if Eqs. (3.39) and (3.40) holds, then Eqs. (3.41) and (3.44) hold. Therefore Eq. (3.42) holds. Conversely, if Eqs. (3.41) and (3.42) hold, then Eqs. (3.40) and (3.43) hold. Therefore Eq. (3.39) hold. $\square$

Definition 3.35. Let $(A, \searrow, \nearrow, \nwarrow, \swarrow)$ be an L-quadri-algebra and $r \in A \otimes A$. Eqs. (3.39)-(3.40) are called *LQ-equation* (a set of equations) in $(A, \searrow, \nearrow, \nwarrow, \swarrow)$. On the other hand, a symmetric bilinear form $\mathcal{B}$ on A is called *2-cocycle* if $\mathcal{B}$ satisfies Eqs. (3.41)-(3.42).

Remark 3.36. In [NB], we show that the *LQ-equation* in an L-quadri-algebra is related to a double construction of the nondegenerate (skew-symmetric) invariant bilinear forms on L-quadri-algebras (or equivalently, the double construction of the nondegenerate (skew-symmetric)

2-cocycles on L-dendriform algebras). In this sense, the LQ -equation in an L-quadri-algebra can be regarded as an analogue of the CYBE in an L-quadri-algebra.

Remark 3.37. Let $(A, \searrow, \nearrow, \nwarrow, \swarrow)$ be an L-quadri-algebra. Then there exists an L-quadri-algebra structure on $A \oplus \mathbb{F}c$ given by (for any $x, y \in A$)

$$(3.46) \quad \begin{aligned} x \bar{\searrow} y &= x \searrow y - \mathcal{B}(y, x)c, & x \bar{\nwarrow} y &= x \nwarrow y - \mathcal{B}(x, y)c, \\ x \bar{\nearrow} y &= x \nearrow y + \mathcal{B}(y, x)c, & x \bar{\swarrow} y &= x \swarrow y + \mathcal{B}(x, y)c, \\ c \bar{\searrow} x &= x \bar{\searrow} c = c \bar{\searrow} c = c \bar{\nwarrow} x = x \bar{\nwarrow} c = c \bar{\nwarrow} c = x \bar{\nearrow} c = c \bar{\nearrow} x = c \bar{\nearrow} c = x \bar{\swarrow} c = c \bar{\swarrow} x = c \bar{\swarrow} c = 0, \end{aligned}$$

if and only if $\mathcal{B}$ satisfies (for any $x, y, z \in A$)

$$(3.47) \quad \mathcal{B}(x \nearrow y, z) = -\mathcal{B}(y, z \vee x - x \wedge z) - \mathcal{B}(z \succ y, x),$$

$$(3.48) \quad \mathcal{B}(x \swarrow y, z) = -\mathcal{B}(z \triangleright x + z \triangleleft x, y) - \mathcal{B}(x, z \vee y).$$

Furthermore, if a bilinear form $\mathcal{B}$ on an L-quadri-algebra $(A, \searrow, \nearrow, \nwarrow, \swarrow)$ satisfying Eqs. (3.47) and (3.48), then it is easy to show that

$$(3.49) \quad \omega(x, y) = \mathcal{B}(x, y) - \mathcal{B}(y, x), \quad \forall x, y \in A,$$

is a 2-cocycle of both the associated vertical L-dendriform algebra $(A, \succ, \triangleleft)$ and the associated depth L-dendriform algebra $(A, \vee, \wedge)$. Obviously, if $\mathcal{B}$ is symmetric, then Eqs. (3.41)-(3.42) and (3.47)-(3.48) coincide.

The following conclusion is obvious.

Proposition 3.38. *Let $(A, \searrow, \nearrow, \nwarrow, \swarrow)$ be an L-quadri-algebra. A skew-symmetric (symmetric respectively) bilinear form $\mathcal{B}$ is invariant (a 2-cocycle respectively) if and only if $\mathcal{B}$ is invariant (a 2-cocycle respectively) on its transpose and symmetries.*

APPENDIX: L-OCTO-ALGEBRAS

Definition A.1. Let A be a vector space with eight bilinear products denoted by

$$\searrow_1, \searrow_2, \nearrow_1, \nearrow_2, \nwarrow_1, \nwarrow_2, \swarrow_1, \swarrow_2: A \otimes A \rightarrow A.$$

$(A, \searrow_1, \searrow_2, \nearrow_1, \nearrow_2, \nwarrow_1, \nwarrow_2, \swarrow_1, \swarrow_2)$ is called an *L-octo-algebra* if (for any $x, y, z \in A$)

$$(A.1) \quad x \searrow_2 (y \searrow_2 z) - (x \ast_{12} y) \searrow_2 z = y \searrow_2 (x \searrow_2 z) - (y \ast_{12} x) \searrow_2 z,$$

$$(A.2) \quad x \searrow_2 (y \nearrow_2 z) - (x \vee_{12} y) \nearrow_2 z = y \nearrow_2 (x \succ_2 z) - (y \wedge_{12} x) \nearrow_2 z,$$

$$(A.3) \quad x \searrow_2 (y \nearrow_1 z) - (x \vee_2 y) \nearrow_1 z = y \nearrow_1 (x \succ_{12} z) - (y \wedge_1 x) \nearrow_1 z,$$

$$(A.4) \quad x \nearrow_2 (y \succ_1 z) - (x \wedge_2 y) \nearrow_1 z = y \searrow_1 (x \nearrow_{12} z) - (y \vee_1 x) \nearrow_1 z,$$

$$(A.5) \quad x \searrow_2 (y \searrow_1 z) - (x *_2 y) \searrow_1 z = y \searrow_1 (x \searrow_{12} z) - (y *_1 x) \searrow_1 z,$$

$$(A.6) \quad x \searrow_2 (y \nwarrow_1 z) - (x \searrow_2 y) \nwarrow_1 z = y \nwarrow_1 (x *_2 z) - (y \nwarrow_1 x) \nwarrow_1 z,$$

$$(A.7) \quad x \nearrow_2 (y \prec_1 z) - (x \nearrow_2 y) \nwarrow_1 z = y \swarrow_1 (x \wedge_{12} z) - (y \swarrow_1 x) \nwarrow_1 z,$$

$$(A.8) \quad x \nearrow_1 (y \prec_{12} z) - (x \nearrow_1 y) \nwarrow_1 z = y \swarrow_2 (x \wedge_1 z) - (y \swarrow_2 x) \nwarrow_1 z,$$

$$(A.9) \quad x \searrow_1 (y \swarrow_{12} z) - (x \succ_1 y) \nwarrow_1 z = y \swarrow_2 (x \vee_1 z) - (y \prec_2 x) \nwarrow_1 z,$$

$$(A.10) \quad x \searrow_1 (y \nwarrow_{12} z) - (x \searrow_1 y) \nwarrow_1 z = y \nwarrow_2 (x *_1 z) - (y \nwarrow_2 x) \nwarrow_1 z,$$

$$(A.11) \quad x \searrow_2 (y \swarrow_1 z) - (x \succ_2 y) \swarrow_1 z = y \swarrow_1 (x \vee_{12} z) - (y \prec_1 x) \swarrow_1 z,$$

$$(A.12) \quad x \searrow_2 (y \swarrow_2 z) - (x \succ_{12} y) \swarrow_2 z = y \swarrow_2 (x \vee_2 z) - (y \prec_{12} x) \swarrow_2 z,$$

$$(A.13) \quad x \nearrow_2 (y \prec_2 z) - (x \nearrow_{12} y) \nwarrow_2 z = y \swarrow_2 (x \wedge_2 z) - (y \swarrow_{12} x) \nwarrow_2 z,$$

$$(A.14) \quad x \searrow_2 (y \nwarrow_2 z) - (x \searrow_{12} y) \nwarrow_2 z = y \nwarrow_2 (x *_2 z) - (y \nwarrow_{12} x) \nwarrow_2 z,$$

where

$$(A.15) \quad x \vee_1 y = x \searrow_1 y + x \swarrow_1 y, \quad x \wedge_1 y = x \nearrow_1 y + x \nwarrow_1 y,$$

$$(A.16) \quad x \vee_2 y = x \searrow_2 y + x \swarrow_2 y, \quad x \wedge_2 y = x \nearrow_2 y + x \nwarrow_2 y,$$

$$(A.17) \quad x \succ_1 y = x \searrow_1 y + x \nearrow_1 y, \quad x \prec_1 y = x \nwarrow_1 y + x \swarrow_1 y,$$

$$(A.18) \quad x \succ_2 y = x \searrow_2 y + x \nearrow_2 y, \quad x \prec_2 y = x \nwarrow_2 y + x \swarrow_2 y,$$

$$(A.19) \quad x \nearrow_{12} y = x \nearrow_1 y + x \nearrow_2 y, \quad x \searrow_{12} y = x \searrow_1 y + x \searrow_2 y,$$

$$(A.20) \quad x \swarrow_{12} y = x \swarrow_1 y + x \swarrow_2 y, \quad x \nwarrow_{12} y = x \nwarrow_1 y + x \nwarrow_2 y.$$

$$(A.21) \quad x \vee_{12} y = x \vee_1 y + x \vee_2 y = x \searrow_{12} y + x \swarrow_{12} y, \quad x \wedge_{12} y = x \wedge_1 y + x \wedge_2 y = x \nearrow_{12} y + x \nwarrow_{12} y,$$

$$(A.22) \quad x \succ_{12} y = x \succ_1 y + x \succ_2 y = x \nearrow_{12} y + x \searrow_{12} y, \quad x \prec_{12} y = x \prec_1 y + x \prec_2 y = x \nwarrow_{12} y + x \swarrow_{12} y,$$

$$(A.23) \quad x *_1 y = x \searrow_1 y + x \nwarrow_1 y + x \nearrow_1 y + x \swarrow_1 y, \quad x *_2 y = x \searrow_2 y + x \nwarrow_2 y + x \nearrow_2 y + x \swarrow_2 y,$$

$$(A.24) \quad x *_2 y = x *_1 y + x *_2 y = x \searrow_{12} y + x \nwarrow_{12} y + x \nearrow_{12} y + x \swarrow_{12} y.$$

Proposition A.2. *Let $(A, \searrow_1, \searrow_2, \nearrow_1, \nearrow_2, \nwarrow_1, \nwarrow_2, \swarrow_1, \swarrow_2)$ be an L -octo-algebra.*

(1) *$(A, \vee_2, \wedge_2, \wedge_1, \vee_1)$ is an L -quadri-algebra.*

(2) *$(A, \succ_2, \succ_1, \prec_1, \prec_2)$ is an L -quadri-algebra.*

(3) $(A, \searrow_{12}, \nearrow_{12}, \nwarrow_{12}, \swarrow_{12})$ is an L -quadri-algebra.

(4) If we define (for any $x, y \in A$)

$$(A.25) \quad x \triangleright_1^2 y = x \searrow_2 y - y \nwarrow_1 x, \quad x \triangleleft_1^2 y = x \nearrow_2 y - y \swarrow_1 x,$$

$$(A.26) \quad x \triangleright_2^1 y = x \searrow_1 y - y \nwarrow_2 x, \quad x \triangleleft_2^1 y = x \nearrow_1 y - y \swarrow_2 x,$$

then $(A, \triangleright_1^2, \triangleleft_1^2, \triangleleft_2^1, \triangleright_2^1)$ is an L -quadri-algebra.

ACKNOWLEDGEMENTS

This work was supported in part by the NSFC (10621101, 10921061), NKBRPC (2006CB805905) and SRFDP (200800550015).

REFERENCES

- [AL] M. Aguiar, J.-L. Loday, Quadri-algebras, J. Pure Appl. Algebra 191 (2004) 205-221.
- [Bai1] C.M. Bai, A unified algebraic approach to the classical Yang-Baxter equation, J. Phys. A: Math. Theor. 40 (2007) 11073-11082.
- [Bai2] C. M. Bai, Left-symmetric bialgebras and an analogy of the classical Yang-Baxter equation, Comm. Contemp. Math. 10 (2008) 221-260.
- [Bai3] C.M. Bai, $\mathcal{O}$ -operators of Loday algebras and analogues of the classical Yang-Baxter equation, Comm. Algebra 38 (2010) 4277-4321.
- [BLN] C.M. Bai, L.G. Liu, X. Ni, Some results on L-dendriform algebras, J. Geom. Phys. 60 (2010) 940-950.
- [Bax] G. Baxter, An analytic problem whose solution follows a simple algebraic identity, Pacific J. Math. 10 (1960), 731-742.
- [Bu] D. Burde, Left-symmetric algebras and pre-Lie algebras in geometry and physics, Cent. Euro. J. Math. 4 (2006) 323-357.
- [Cha] F. Chapoton, Un théorème de Cartier-Milnor-Moore-Quillen pour les bigèbres dendriformes et les algèbres braces, J. Pure Appl. Algebra 168 (2002) 1-18.
- [CP] V. Chari, A. Pressley, A guide to quantum groups, Cambridge University Press, Cambridge (1994).
- [Chu] B.Y. Chu, Symplectic homogeneous spaces, Trans. Amer. Math. Soc. 197 (1974) 145-159.
- [E] K. Ebrahimi-Fard, Loday-type algebras and the Rota-Baxter relation, Lett. Math. Phys. 61 (2002) 139-147.
- [EG] K. Ebrahimi-Fard, L. Guo, On products and duality of binary, quadratic, regular operads, J. Pure Appl. Algebra 200 (2005) 293-317.
- [EMP] K. Ebrahimi-Fard, D. Manchon, F. Patras, New identities in dendriform algebras, J. Algebra 320 (2008) 708-727.
- [Fo] L. Foissy, Les algèbres de Hopf des arbres enracinés décorés II, Bull. Sci. Math. 126 (2002) 249-288.
- [Fr1] A. Frabetti, Dialgebra homology of associative algebras, C. R. Acad. Sci. Paris 325 (1997) 135-140.
- [Fr2] A. Frabetti, Leibniz homology of dialgebras of matrices, J. Pure Appl. Algebra 129 (1998) 123-141.
- [G] M. Gerstenhaber, The cohomology structure of an associative ring, Ann. Math. 78 (1963) 267-288.
- [GS] I.Z. Golubschik, V.V. Sokolov, Generalized operator Yang-Baxter equations, integrable ODEs and nonassociative algebras, J. Nonlinear Math. Phys. 7 (2000) 184-197.
- [Ko] J.-L. Koszul, Domaines bornés homogènes et orbites de groupes de transformation affines, Bull. Soc. Math. France 89 (1961) 515-533.
- [Ku] B.A. Kupersmidt, What a classical r -matrix really is, J. Nonlinear Math. Phys. 6 (1999) 448-488.
- [Le1] P. Leroux, Ennea-algebras, J. Algebra 281 (2004) 287-302.
- [Le2] P. Leroux, Construction of Nijenhuis operators and dendriform trialgebras, Int. J. Math. Sci. (2004) 2595-2615.
- [Le3] P. Leroux, On some remarkable operads constructed from Baxter operators, arXiv: math.QA/0311214.
- [Lo1] J.-L. Loday, Dialgebras, in Dialgebras and related operads, Lecture Notes in Math. 1763 (2002) 7-66.
- [Lo2] J.-L. Loday, Arithmetree, J. Algebra 258 (2002) 275-309.
- [Lo3] J.-L. Loday, Scindement d'associativité et algèbres de Hopf, Proceedings of the Conference in honor of Jean Leray, Nantes (2002), Séminaire et Congrès (SMF) 9 (2004) 155-172.
- [LR1] J.-L. Loday, M. Ronco, Hopf algebra of the planar binary trees, Adv. Math. 139 (1998) 293-309.

- [LR2] J.-L. Loday, M. Ronco, Algèbre de Hopf colibres, C.R. Acad. Sci. Paris 337 (2003) 153-158.
- [LR3] J.-L. Loday, M. Ronco, Trialgebras and families of polytopes, in “Homotopy Theory: Relations with Algebraic Geometry, Group Cohomology, and Algebraic K-theory”, Contemp. Math. 346 (2004) 369-398.
- [NB] X. Ni, C.M. Bai, L-dendriform bialgebras and L-quadri-bialgebras, preprint, 2009.
- [Ron] M. Ronco, Primitive elements in a free dendriform algebra, Contemp. Math. 267 (2000) 245-263.
- [Rot] G.-C. Rota, Baxter algebras and combinatorial identities I, Bull. Amer. Math. Soc. 75 (1969) 325-329.
- [Sc] R. Schafer, An introduction to nonassociative algebras, Dover Publications Inc., New York (1995).
- [Se] M.A. Semenov-Tian-Shansky, What is a classical R-matrix? Funct. Anal. Appl. 17 (1983) 259-272.
- [Sh] H. Shima, Homogeneous Hessian manifolds, Ann. Inst. Fourier, Grenoble 30 (1980) 91-128.
- [Va] B. Vallette, Manin products, Koszul duality, Loday algebras and Deligne conjecture, J. Reine Angew. Math. 620 (2008) 105-164.
- [Vi] E.B. Vinberg, Convex homogeneous cones, Transl. of Moscow Math. Soc. 12 (1963) 340-403.

CHERN INSTITUTE OF MATHEMATICS & LPMC, NANKAI UNIVERSITY, TIANJIN 300071, P.R. CHINA
E-mail address: liuligong@mail.nankai.edu.cn

CHERN INSTITUTE OF MATHEMATICS & LPMC, NANKAI UNIVERSITY, TIANJIN 300071, P.R. CHINA
E-mail address: xiangn_math@yahoo.cn

CHERN INSTITUTE OF MATHEMATICS & LPMC, NANKAI UNIVERSITY, TIANJIN 300071, P.R. CHINA
E-mail address: baicm@nankai.edu.cn